

Gulf of Mexico Harmful Algal Bloom Bulletin


19 August 2004 National Ocean Service/NCCOS and CSC NESDIS/CoastWatch and NDBC Last bulletin: August 2, 2004

Analysis SW Florida:


Algal blooms developed from Venice to Naples in response to Hurricane Charley. Diatom blooms are common after hurricanes. Generally low to slightly moderate probability of Karenia brevis in the bloom area from Boca Grande to Venice. Low probability from Sanibel south. No development anticipated over next several days.

--Stumpf

Please note the following restrictions on all SeaWiFS imagery derived from CoastWatch.



Chlorophyll concentration from satellite with possible HAB areas shown by red polygon(s). Cell concentration sampling data from August 15, 2004 shown as red squares (high), red triangles (medium), red diamonds (low b), red circles (low a), orange circles (very low b), yellow circles (very low a), green circles (present), and black "X" (not present).



Wind speed and direction are averaged over 12 hours from measurements made on buoys. Length of line indicates speed; angle indicates direction. Red indicates that the wind direction favors upwelling near the coast. Values to the left of the dotted vertical line are measured values; values to the right are forecasts.


Variable forecast winds

^{1.} These data are restricted to civil marine applications only; i.e. federal, state, and local government use/distribution is permitted .

^{2.} Distribution for military, or commercial purposes is NOT permitted.

^{3.} There are restrictions on Internet/Web/public posting of these data.

Image products may be published in newspapers. Any other publishing arrangements must receive OrbImage approval via the CoastWatch Program.



Blooms shown in red (see p. 1 analysis and image for interpretation)

Wind conditions from Egmont Key, FL

